


**Cancer cell's neuroendocrine feature can be acquired through cell-cell fusion during cancer-neural stem cell interaction**

Liyuan Yin<sup>1,2</sup>, Peizhen Hu<sup>2</sup>, Xianping Shi<sup>3</sup>, Weiping Qian<sup>4</sup>, Haiyen E. Zhau<sup>2</sup>, Leland W. K. Chung<sup>2</sup>, Michael S. Lewis<sup>5</sup>, Stephen J. Pandol<sup>3</sup>, and Ruoxiang Wang<sup>2,5</sup>

<sup>1</sup>Lung Cancer Center, West China Hospital, Sichuan University, Chengdu, China; <sup>2</sup>Uro-Oncology Research, <sup>3</sup>Department of Medicine, Cedars-Sinai Medical Center, Los Angeles, CA, USA; <sup>5</sup>Department of Pathology, Greater Los Angeles Veterans Affairs Health System, Los Angeles, CA, USA; and <sup>4</sup>Department of Surgery, Emory University School of Medicine, Atlanta, GA, USA.

## Supplementary Data

Figure S1



**Figure S1. Cell number changes during co-culture.** Cells in co-culture were prepared every 7 days in single-cell suspension with trypsin-EDTA treatment; and were counted with a TC10 counter based on cell size. Quadruple counts were obtained from each sample. The data are shown with the mean of quadruple counts. For all the data points, standard deviations are less than 5% of the mean and are not shown.